


Center for  
Molecular  
Modeling

## Computational Materials Physics



Department of  
Materials Science  
and Engineering

### supercells (impurities and surfaces)

Stefaan.Cottenier@ugent.be  
Technologiepark 903, Zwijnaarde

<http://molmod.ugent.be>  
<http://www.ugent.be/ea/dmse/en>  
my talks on Youtube: <http://goo.gl/P2b1Hs>

---

---

---

---

---


---

---


---


### the supercell concept


Put a single atom in an otherwise empty, big unit cell:





→


←→10-15 Å

















fcc Bravais lattice:  
12 "nearest" neighbours =  
as spherically symmetric as possible.

---

---

---

---

---


---

---


---

### the supercell concept

Replicate the chemical unit cell (here 2x2 in 2D) and replace one atom by an impurity:



→



---

---

---


---

---

---

---

---




---

---

---

---

---


---

---

---

---

---




---

---

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---


---

---

**example: Fe(001)**

new convergence criteria to be checked for a physically relevant model :

- slab thickness
- vacuum thickness



---

---

---

---


---

---

---

---

**example: Fe(001)**



---

---

---


---

---

---

---

---



how to model a surface covered by a monolayer of another phase? (=extremely thin coating)

---

---

---


---

---

---

---

---



how to model a surface with an isolated impurity atom attached to it? (=adsorbate)

---

---

---

---

---

---

---

---